



Expression of cytochrome P450 2A13 in human non-small cell lung cancer and its clinical significance

Li Sun[✉], Xiaoli Fan

Department of Histology and Embryology, College of Basic Medicine of Guilin Medical University, Guilin, Guangxi 541004, China.

Received 09 March 2012, Revised 11 June 2012, Accepted 13 January 2013, Epub 24 March 2013

Abstract

Lung cancer is one of the most important causes of cancer-related mortality worldwide. Human cytochrome P450 2A13 enzyme (CYP2A13) is predominantly expressed in the respiratory tract and could catalyze various carcinogens. In this study, we quantified CYP2A13 expression in non-small cell lung cancer (NSCLC) tissues and examined the relation between CYP2A13 and clinicopathologic factors. Thirty-five paired lung cancer and normal tissues were studied for the expression of the *CYP2A13* gene by using real-time PCR and Western blotting assays. We also investigated the relationship between *CYP2A13* expression and clinicopathologic factors such as age, gender, histology and lymph node status in tumor tissues. SPSS (17.0) statistical software was applied for data analysis. The real-time PCR results showed that there was no significant difference in the *CYP2A13* mRNA transcript levels between tumor and paired normal tissues in the 35 samples and in 12 paired squamous cell carcinomas. In adenocarcinoma, the expression of *CYP2A13* mRNA in tumor tissues was 12.5% of that in adjacent tissues ($P < 0.05$) and it was not associated with age, gender, histology and lymph node status of the patients. The amounts of CYP2A13 proteins detected by Western blotting assays correlated well with those of the corresponding mRNAs. In conclusion, the expression of CYP2A13 was downregulated in lung adenocarcinoma. CYP2A13 may be involved in the development and progression of lung adenocarcinoma.

Keywords: cytochrome P450 2A13 (CYP2A13), non-small lung cancer, real-time PCR

INTRODUCTION

Human cytochrome P450 (CYP450) enzymes are the principal enzymes involved in the metabolic activation of many compounds, such as drugs, environmental toxicants and chemical carcinogens^[1]. Variations of P450 enzymes in expression or activity can contribute substantially to inter-individual differences

in therapeutic efficacy or toxicant susceptibility^[2]. *CYP2A13* is a member of the *CYP2A* gene subfamily of P450s, which is predominantly expressed in the nasal mucosa and also in the lung and trachea^[3,4]. *CYP2A13* is highly effective in the metabolic activation of many respiratory-tract toxicants, especially 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK), which is a tobacco-specific carcinogen^[5-7].

This work was supported by the Project sponsored by the Scientific Research Foundation for the Returned Overseas Chinese Scholars, State Education Ministry (No. Z2005-2-45006).

✉Corresponding author: Prof. Li Sun, Department of Histology and

Embryology, College of Basic Medicine of Guilin Medical University, 109 N.2nd Huan Cheng, Guilin, Guangxi 541004, China. Tel:+86-7735895086, E-mail: sunli2196@glmc.edu.cn.

The authors reported no conflict of interests.

Furthermore, it was found that CYP2A13 is active for the metabolism of aflatoxin B1^[8]. The expression of CYP2A13 may also substantially influence the incidence of respiratory tumor. However, the precise role of CYP2A13 in carcinogenesis and tumor progression is still unclear. In the present study, we measured the expressions of CYP2A13 in lung cancer and paired normal tissues and examined their relationship with clinicopathological factors.

MATERIALS AND METHODS

Tissue samples

Thirty-five surgical specimens of adenocarcinoma and squamous cell carcinoma were used for the current study. No patients had received chemotherapy before surgery. These patients underwent surgery between 2009 and 2010 at Guilin Medical University Hospital, Guilin, Guangxi, China. Tumor and normal lung tissue samples were obtained from the resected specimens, and snap-frozen in liquid nitrogen and stored at -80°C for further analysis. The study protocol was approved by the local institutional board at the authors' affiliated institution. The use of human tissue specimens for experimental purposes was done in accordance with the established institutional and national guidelines regarding experimental use of human tissues.

Chemicals and reagents

PrimeScript RT Reagent Kit with gDNA Eraser (Perfect Real Time) and SuperReal PreMix (SYBR Green) were purchased from Takara (Dalian, Shandong, China) and Tiangen (Beijing, China), respectively. Cell lysis buffer for Western blotting assays and IP and Enhanced BCA Protein Assay Kit BCA was purchased from Beyotime (Beijing, China).

Preparation of antibody

Rabbit polyclonal antibody against human CYP2A13 and biotinylated goat anti-rabbit IgG were obtained from Bioworld (Louis Park, MN, USA). Mouse monoclonal antibody against human β -actin goat anti-mouse IgG were purchased from Beyotime (Beijing, China).

Quantitative analysis of CYP2A13 mRNA

Total RNA (1 μ g) was extracted from frozen samples by using TRIZOL reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocol. The quality of the RNA samples was determined by electrophoresis through 1% agarose gels and the 18S and 28S bands were visualized under ultraviolet light. RNA samples were cleaned by using PrimeScript RT

Reagent Kit with gDNA Eraser to ensure that no contaminating genomic DNA was present and then converted to cDNA (Takara). Gene expression was measured by SYBR green real-time PCR by using ABI 7500 FAST detection system applied biosystems (ABI, New York, USA). β -actin, a housekeeping gene, was used as an internal control. The primer sequences were as follows: CYP2A13-forward: 5'-ATA AGA AGG GGC AGT TTA AGA AGA G-3', reverse: 5'-ATG ATG GTG GTG AAG AAG AGA AAG-3', β -actin-forward: 5'-AGC GAG CAT CCC CCA AAG TT-3', reverse: 5'-GGG CAC GAA GGC TCA TCA TT-3'. The PCR system was performed in 20 μ L buffers containing 1 μ L cDNA, 0.4 μ L of 50 \times ROX Reference Dye and 50 pmol of primer pairs (Tiangen). Thermocycling conditions consisted of a pre-incubation for 15 minutes at 95°C and 60°C for 30 seconds, followed by 40 cycles of denaturation for 10 seconds at 94°C. PCR products were separated by 2% agarose gel electrophoresis and sequenced to identify the specificity.

The experiment used comparative CT method to present relative gene expression (also known as the $2^{-\Delta\Delta C_t}$ method). The PCR efficiency of the CYP2A13 gene must be similar to the internal control gene β -actin. The PCR efficiency of CYP2A13 and β -actin were included in the equation; therefore, the differences in the efficiency between target and internal control will be accounted for in the calculation. The equation is: $2^{-\Delta\Delta C_t} = [CT_{CYP2A13} - CT_{\beta-actin}]_{tumor} - [CT_{CYP2A13} - CT_{\beta-actin}]_{adjacent\ normal\ tissue}$. In this way, the data may be interpreted as "the expression of the CYP2A13 gene to β -actin in the lung cancer compared with the paired normal tissue".

Western blotting assays

The proteins prepared from tissue samples were subjected to SDS-polyacrylamide gel (12%), and then transferred electrophoretically to a nitrocellulose membrane. The membrane was blocked in 5% non-fat dry milk for 1 hour at 37°C before incubation with the monoclonal anti-human CYP2A13 antibody against human and β -actin at room temperature for 2 hours. After washing with tris buffered saline, the membrane was incubated with anti-rabbit IgG and anti-mouse IgG (both conjugated with horseradish peroxidase) for 1 hour at room temperature to detect CYP2A13 and β -actin. The immunoblots were visualized by enhanced chemiluminescence (ECL) in the manufacturer's protocol.

Statistical analysis

For statistical analyses, SPSS 17.0 statistical software (SPSS Inc., Chicago, IL, USA) was performed.

Table 1 Patient demographic and baseline characteristics

Variables	Patients [n(%)]
Age	
≤ 55 years	15 (42.8)
> 55 years	20 (57.2)
Gender	
Male	24 (68.6)
Female	11 (31.7)
Lymph node status	
Node-negative	16 (45.7)
Node-positive	19 (54.3)
Histology	
Squamous cell carcinomas	23 (65.7)
Adenocarcinomas	12 (34.3)

The expressions of CYP2A13 between lung cancer and paired normal tissue were investigated using the paired-sample *t*-test and the relationship between CYP2A13 expression and clinicopathological factors were examined by using independent-sample *t* test. *P* value < 0.05 was considered significant.

RESULTS

Patient demographic and baseline characteristics are

shown in **Table 1**. We used H&E staining to identify different pathological types (**Fig. 1**). Twelve (34.3%, 12/35) patients had adenocarcinoma and 23 (65.7%) patients had squamous cell carcinoma. Nineteen (54.3%) patients had metastasis to lymph nodes.

CYP2A13 mRNA levels in adult human lung tissue samples

The *CYP2A13* mRNA transcript levels were 11.92 ± 4.08 for the tumor tissues ($n = 35$) and 10.18 ± 5.88 for adjacent paired normal tissues ($n = 35$) (**Table 2**). Paired-sample *t*-test was used to analyze the expression levels of *CYP2A13* mRNA in the 35 breast cancer samples and paired normal tissues and no significant difference was found between them ($P = 0.059$). The *CYP2A13* mRNA transcript levels were 11.53 ± 4.45 for the squamous cell carcinoma tissues ($n = 12$) and 12.31 ± 6.76 for adjacent paired normal tissues ($n = 12$). The expression level of *CYP2A13* mRNA transcripts between squamous cell carcinoma and matched normal tissues showed no significant difference. ($P = 0.604$). Furthermore, lower expression of *CYP2A13* mRNA in adenocarcinoma was observed (12.12 ± 3.95) compared with that in adjacent nor-




Fig. 1 H&E staining of lung cancer tissues (100×). A: adenocarcinomas. Slice shows gland forming tumor that grows into papillary pattern. B: paired normal tissue of adenocarcinomas; C: squamous cell carcinomas. Tumor cells show well formed keratin pearl. D: paired normal tissue of squamous cell carcinomas.

Table 2 CYP2A13 mRNA expression in non-small cell lung cancer (NSCLC) tissues and adjacent normal tissues

Variables	<i>n</i>	CYP2A13 mRNA levels	$2^{-\Delta\Delta CT}$	<i>P</i>
Total NSCLC	35	11.92 ± 4.08		
Paired adjacent normal tissue	35	10.18 ± 5.88		0.059
Tumor types				
Squamous cell carcinoma	12	11.53 ± 4.45		
Paired adjacent normal tissue	12	12.31 ± 6.76		0.604
Adenocarcinoma	23	12.12 ± 3.95		
Paired adjacent normal tissue	23	9.07 ± 5.18	0.125	0.008

mal tissues (9.07 ± 5.18) with a statistical difference between the two groups ($P = 0.008$), suggesting that CYP2A13 mRNA was downregulated in lung adenocarcinoma tissues compared to normal tissues.

CYP2A13 mRNA expression and clinicopathological factors

We examined the relationship between the level of CYP2A13 mRNA and clinicopathological factors in tumors by using independent-sample t test. The expression level of CYP2A13 mRNA in tumor tissues was not associated with age, gender, histology and lymph node status of patients (**Table 3**).

CYP2A13 protein expression in lung tissue

The expression of CYP2A13 protein was analyzed by Western blotting assays. We randomly chose four representative cases (**Fig. 2A**). CYP2A13 proteins expressed in the tumor tissues were compared with those of the adjacent normal tissues. We found that CYP2A13 protein levels corresponded to its mRNA transcript levels (**Fig. 2B**). The expressions of CYP2A13 proteins were consistent with variations in the levels of the corresponding CYP2A13 mRNA transcripts in the same tissue samples, and the ratio of CYP2A13 protein in tumor: normal tissue almost correlated with that of mRNA (**Fig. 2C**).

DISCUSSION

The cytochrome P450 superfamily are the major enzymes involved in drug metabolism and bioactivation, and CYP catalyzes various environmental protoxicants and chemical procarcinogens, such as 4-aminobiphenyl, naphthalene, styrene and toluene^[9,10]. There are two functional members of the human CYP2A gene subfamily: CYP2A6 and CYP2A13. CYP2A6 is an efficient human enzyme in the metabolism of coumarin and nicotine and it is mainly expressed in human liver, whereas CYP2A13 is predominantly expressed in the respiratory tract. The cDNA and protein sequences of CYP2A13 are highly similar to those of CYP2A6 with only 4.7% and 6.5% difference in the nucleotide coding sequence and amino acid sequence, respectively^[11]. CYP2A13 is much more active for nicotine, cotinine, and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone metabolisms and especially alphatoxin B1. Recently, it is found that CYP2A13 catalyzes the metabolic activation of NNK through α -hydroxylation^[12]. NNK and aflatoxin B1 are all potential carcinogens, and it has been reported previously that CYP2A13 was associated with lung adenocarcinomas risk^[3]. For CYP2A13, variations in its expression level or metabolic capacity may significantly alter the extent of

Table 3 Relationship between CYP2A13 mRNA expression and clinicopathological factors

Features	No. of patients	CYP2A13 mRNA	<i>P</i>
Age			
≤ 55 years	15	12.06 ± 4.44	
> 55 years	20	11.93 ± 3.90	0.794
Gender			
Male	24	11.08 ± 3.40	
Female	11	13.74 ± 4.97	0.072
Lymph node status			
Node-negative	16	12.26 ± 4.50	
Node-positive	19	11.73 ± 3.46	0.724
Histology			
Squamous cell carcinomas	12	11.53 ± 4.46	
Adenocarcinomas	23	12.12 ± 3.96	0.694




Fig. 2 Expression of CYP2A13 in lung cancer tissues. A: Western blotting for detecting the expression of CYP2A13 in the samples (T: tumor; N: adjacent normal tissue). B: The expression analysis of CYP2A13 at mRNA and protein levels. C: The comparison of CYP2A13 between tumor and paired normal tissues. The expression of the CYP2A13 proteins were consistent with variations in the levels of the corresponding *CYP2A13* mRNAs in the same tissue samples.

metabolic activation of NNK and other xenobiotic substrates in the respiratory tract, which may lead to altered susceptibility to tobacco-related tumorigenesis. Wang et al.^[13] studied the relationship between CYP2A13 genetic polymorphisms and the susceptibility of lung cancer. The results showed that genetic polymorphisms of CYP2A13 may have a significant impact on human susceptibility to lung cancers, especially adenocarcinomas.

In this study, we provided the first quantification of the amount of *CYP2A13* mRNA between lung cancer tissues and its adjacent normal tissues by using quantitative RT-PCR. The expression of CYP2A13 levels in lung adenocarcinoma tissue was significantly lower than that in normal tissue. Previously, it is found that in the hepatic cancer, many P450s are downregulated, and the expression obviously decreased in tumors compared with the matched normal tissue^[14]. The expression level of CYP2A13 in tumor was not associated with age, gender, histological differentiation and lymph node status of patients. Etzel et al.^[15] reported that many cytochrome P450 (CYP) enzymes were involved in estrogen metabolism and regulated by estrogens. They thought that these genes may be relevant to gender-specific differences in lung cancer risk in early-onset lung cancer, where a high proportion of women were observed. A case-control study

of Timofeeva et al.^[16] demonstrated that *CYP1B1* and *CYP2A13* genotypes may be related to individual susceptibility to early-onset lung cancer in women. The expression of P450s may be markers of prognosis or influence the metabolism of many xenobiotics by P450s^[17,18].

As the expression of mRNA may not truly reflect the level of protein, we detected the level of protein by using Western blotting assays. The results showed that the levels of CYP2A13 proteins varied with the levels of the corresponding *CYP2A13* mRNAs in the same sample. This result is consistent with the observation of Zhang et al.^[5]. So far, two research groups had used immunohistochemistry to analyze intracellular distribution of CYP2A13 and they generated specific antibody against CYP2A13. They found that CYP2A13 was present in normal trachea but not in peripheral lung tissues. Thus, it demonstrated that the expression of CYP2A13 is not uniform in the lung, which supported the negative results of immunoblotting analysis of the microsomes of the lung tissues. Zhu^[19] reported that CYP2A13 protein was not detected in any large cell carcinomas ($n = 3$), adenocarcinomas ($n = 6$), squamous carcinomas ($n = 3$), and so on, which was opposite to the result of Fukami^[20], who found that CYP2A13 was increased in NSCLC. As the reason for this discrepancy is not clear, it may partly depend

on the different reactivities of the antibodies, samples or individual differences. Williams and Sandler^[21] reported that its high expression in the respiratory tract and effective metabolism of NNK are consistent with the observation that most smoking-related lung cancers are bronchogenic.

CYP2A13 metabolizes many carcinogenic components of tobacco smoke, such as NNK. The mechanisms may be different between smokers and nonsmokers. The number of samples in this study is limited, and the smoking status of the study population was unknown. In summary, we showed that the expression of CYP2A13 was lower in adenocarcinoma tissues compared with paired normal tissue. It is interesting that CYP2A13 expression was downregulated in adenocarcinoma, which suggests that CYP2A13 may play an important role in the development and progression of lung adenocarcinoma.

References

- [1] Guengerich FP. Cytochrome P450: what have we learned and what are the future issues. *Drug Metab Rev* 2004; 36: 159-97.
- [2] Ingelman-Sundberg M, Sim SC, Gomez A, Rodriguez-Antona C. Influence of cytochrome P450 polymorphisms on drug therapies: pharmacogenetic, pharmacoeconomic and clinical aspects. *Pharmacol Ther* 2007; 116: 496-526.
- [3] Su T, Bao ZP, Zhang QY, Smith TJ, Hong JY, Ding X. Human cytochrome P450 CYP2A13: Predominant expression in the respiratory tract and its high efficiency metabolic activation of a tobacco-specific carcinogen, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Cancer Res* 2000; 60: 5074-9.
- [4] Zhu LR, Thomas PE, Lu G, Reuhl KR, Yang GY, Wang LD, et al. CYP2A13 in human respiratory tissues and lung cancers: an immunohistochemical study with a new peptide-specific antibody. *Drug Metab Dispos* 2006; 34: 1672-6.
- [5] Zhang X, D'Agostino J, Wu H, Zhang QY, von Weymarn L, Murphy SE, et al. CYP2A13: variable expression and role in human lung microsomal metabolic activation of the tobacco-specific carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *J Pharmacol Exp Ther* 2007; 323: 570-8.
- [6] Ding X, Kaminsky LS. Human extrahepatic cytochromes P450: Function in xenobiotic metabolism and tissue-selective chemical toxicity in the respiratory and gastrointestinal tracts. *Annu Rev Pharmacol Toxicol* 2003; 43: 149-73.
- [7] Jalas JR, Hecht SS, Murphy SE. Cytochrome P450 enzymes as catalysts of metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone, a tobacco specific carcinogen. *Chem Res Toxicol* 2005; 18: 95-110.
- [8] He XY, Tang LL, Wang SL, Cai QS, Wang JS, Hong JY. Efficient activation of aflatoxin B-1 by cytochrome P450 2A13, an enzyme predominantly expressed in human respiratory tract. *Int J Cancer* 2006; 118: 2665-71.
- [9] Nakajima M, Itoh M, Sakai H, Fukami T, Katoh M, Yamazaki H, et al. CYP2A13 expressed in human bladder metabolically activates 4-aminobiphenyl. *Int J Cancer* 2006; 119: 2520-6.
- [10] Fukami T, Katoh M, Yamazaki H, Yokoi T, Nakajima M. Human cytochrome P450 2A13 efficiently metabolizes chemicals in air pollutants: naphthalene, styrene, and toluene. *Chem Res Toxicol* 2008; 21: 720-5.
- [11] Brian D, Smith J, Sander L, Lushington GH, Stout CD, Scott EE. Structure of the human lung cytochrome P450 2A13. *Biol Chem* 2007; 282: 17306-13.
- [12] Chiang HC, Wang CY, Lee HL, Tsou TC. Metabolic effects of CYP2A6 and CYP2A13 on 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK)-induced gene mutation--a mammalian cell-based mutagenesis approach. *Toxicol Appl Pharmacol* 2011; 253: 145-52.
- [13] Wang H, Tan W, Hao B, Miao X, Zhou G, He F, et al. Substantial reduction in risk of lung adenocarcinoma associated with genetic polymorphism in CYP2A13, the most active cytochrome P450 for the metabolic activation of tobacco-specific carcinogen NNK. *Cancer Res* 2003; 63: 8057-61.
- [14] Liu Xiufeng, Shi Ruihua, Wang Xuehao, Zhang Guoxin, Zhu Hong, Gao Hengjun, et al. Down-regulated Expression of Cytochrome P450 Related Genes Involved in Hepatocellular Carcinoma. *Chinese-German J Clin Oncol* 2006; 5: 159-64.
- [15] Etzel CJ, Lu M, Merriman K, Liu M, Vaporciyan A, Spitz MR. An epidemiologic study of early onset lung cancer. *Lung Cancer* 2006; 52: 129-34.
- [16] Timofeeva MN, Kropp S, Sauter W, Beckmann L, Rosenberger A, Illig T, et al. CYP450 polymorphisms as risk factors for early-onset lung cancer: gender-specific differences. *Carcinogenesis* 2009; 30: 1161-9.
- [17] Downie D, McFadyen MC, Rooney PH, Cruickshank ME, Parkin DE, Miller ID, et al. Profiling cytochrome P450 expression in ovarian cancer: identification of prognostic markers. *Clin Cancer Res* 2005; 11: 7369-75.
- [18] Kumarakulasingham M, Rooney PH, Dundas SR, Telfer C, Melvin WT, Curran S, et al. Cytochromes P450 profile of colorectal cancer: identification of markers of prognosis. *Clin Cancer Res* 2005; 11: 3758-65.
- [19] Zhu LR, Thomas PE, Lu G, Reuhl KR, Yang GY, Wang LD, et al. CYP2A13 in human respiratory tissues and lung cancers: an immunohistochemical study with a new peptide-specific antibody. *Drug Metab Dispos* 2006; 34: 1672-6.
- [20] Fukami T, Nakajima M, Matsumoto I, Zen Y, Oda M, Yokoi T. Immuno-histochemical analysis of CYP2A13 in various types of human lung cancers. *Cancer Sci* 2010; 101: 1024-8.
- [21] Williams MD, Sandler AB. The epidemiology of lung cancer. *Cancer Treat Res* 2001; 105: 31-52.