



Ovalicin attenuates atopic dermatitis symptoms by inhibiting IL-31 signaling and intracellular calcium influx

Sung-Hyun Hwang^{1,2}, Yeseul Yang^{1,2}, Yeji Jeong^{1,2}, Yongbaek Kim^{1,3,✉}

¹Laboratory of Clinical Pathology, College of Veterinary Medicine, Seoul National University, Seoul 08826, Republic of Korea;

²BK21 PLUS Program for Creative Veterinary Science Research, College of Veterinary Medicine, Seoul National University, Seoul 08826, Republic of Korea;

³Research Institute for Veterinary Science, College of Veterinary Medicine, Seoul National University, Seoul 08826, Republic of Korea.

Supplementary Table 1 Sequences of the primers used for Real-time RT-PCT

Gene name	Sequence (5'→3')
<i>IL31RA</i>	Forward: AGG CCT TTG CTG CCA CTT TC
	Reverse: CAG GAA GCA TTC CCA GGC TC
<i>OSMR</i>	Forward: GTA AGG TGG ACC GAG GAG ATG
	Reverse: GAG CAA GCC AGT GTC TCT C
<i>HRH2</i>	Forward: GCA GTC TGA CTA ACT GCT TC
	Reverse: GGC TGA TCA TGA AGA GGT TG
<i>IFNG</i>	Forward: GCG CAA GGC GAT AAA TGA AC
	Reverse: CTG ACT CCT TTT CCG CTT CC
<i>IL4</i>	Forward: TCA CCA GCA CCT TTG TCC AC
	Reverse: CGC TTG TGT TCT TTG GAG CA
<i>TRPV1</i>	Forward: GGC TGT CTT CAT CAT CCT GCT GCT
	Reverse: GTT CTT GCT CTC CTG TGC GAT CTT GT
<i>GAPDH</i>	Forward: ATC ACT GCC ACC CAG AAG AC
	Reverse: TCA GCT CAG GGA TGA CCT TG

✉Corresponding author: Yongbaek Kim, Laboratory of Clinical Pathology, College of Veterinary Medicine, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Republic of Korea. Tel/Fax: +82-2-880-1273/+82-2-873-1213, E-mail: yongbaek@snu.ac.kr.

Received: 22 January 2021; Revised: 09 April 2021; Accepted: 14 April 2021; Published online: 16 June 2021




Fig. 1 Chelation of intracellular calcium by BAPTA-am treatment in LPS-pretreated DH82 cells. Representative image of Fluo-4AM and DAPI staining in LPS-pretreated DH82 cells with/without BAPTA-AM treatment. Scale bar: 400 μ m. LPS: lipopolysaccharide.




Fig. 2 Macrophage infiltration, weight gain and differential WBC counts in experimental mice. A: Representative immunohistochemical staining for IBA1 in skin tissue of DNCB-untreated control and 0.1% DNCB solution-treated AD mice with/without ovalicin treatment. Scale bar: 200 μ m. B: Mean weight of DNCB-untreated control and 0.1% DNCB solution-treated AD mice with/without ovalicin treatment. Data are presented as mean \pm SEM ($n=5$). C: Percentage of differential WBC counts, including neutrophils, lymphocytes, monocytes, and eosinophils in DNCB-untreated control and 0.1% DNCB solution-treated AD mice with/without ovalicin treatment. Error bars indicate SEM ($n=5$). ** $P<0.01$ and *** $P<0.001$ vs. DNCB-untreated control mice without ovalicin treatment. ## $P<0.01$ vs. DNCB-treated mice without ovalicin treatment. Statistical analysis was conducted by ANOVA and Tukey's test (pairwise) for comparison with samples. WBC: white blood cell; DNCB: 2,4-dinitrochlorobenzene.